

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

2019 Workshop: Interviewers and Their Effects
from a Total Survey Error Perspective

Sociology, Department of

2020

Chapter 7: Statistical Identification of Fraudulent Interviews in Surveys: Improving Interviewer Controls Appendix 7

Silvia Schwanhäuser

Joseph W. Sakshaug

Yuliya Kosyakova

Frauke Kreuter

Follow this and additional works at: <https://digitalcommons.unl.edu/sociw>



Part of the [Quantitative, Qualitative, Comparative, and Historical Methodologies Commons](#)

This Article is brought to you for free and open access by the Sociology, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in 2019 Workshop: Interviewers and Their Effects from a Total Survey Error Perspective by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Chapter 7: Statistical Identification of Fraudulent Interviews in Surveys: Improving
Interviewer Controls

Appendix 7

Silvia Schwanhäuser, Joseph W. Sakshaug, Yuliya Kosyakova, Frauke Kreuter

Full book reference:

Olson, K., J. D. Smyth, J. Dykema, A. L. Holbrook, F. Kreuter, B. T. West. 2020. *Interviewer Effects from a Total Survey Error Perspective*. Boca Raton: CRC Press.

Appendix 7

Table A7A.1 Number of identical response patterns

Pattern occurred	Frequency of patterns	Number of interviews
1 time	4,067	4,067
2 times	91	182
3 times	19	57
4 times	15	60
5 times	15	75
6 times	8	48
7 times	4	28
8 times	1	8
9 times	3	27
10 times	2	20
11 times	1	11
13 times	1	13
15 times	1	15
16 times	1	16
20 times	1	20
169 times	1	169
Total number of interviews		4,816

Source: IAB-BAMF-SOEP Survey of Refugees in Germany, 2016, own calculations.

Explanation: The column “pattern occurred” shows how often a specific response pattern occurred across different interviews. The column “frequency of pattern” indicates how many patterns occurred with the specific repetition number of column “pattern occurred”. The column “number of interviews” shows how many interviews were affected by this. For example, two patterns (10-10-10-10-10-10-10-10-0 and 10-10-10-10-8-10-10-10-10) occurred 10 times i.e. in 10 interviews resulting in 20 interviews in total.